

SUSY breaking in S^1/Z_2 orbifold models*

M. Chaichian^{a †}, A. Kobakhidze^{a,c ‡} and M. Tsulaia^{b,c §}

^a*HEP Division, Department of Physics, University of Helsinki
and Helsinki Institute of Physics,
Helsinki, 00014, Finland*

^b*Bogoliubov Laboratory of Theoretical Physics, JINR
Dubna, 141980, Russia*

^c*The Andronikashvili Institute of Physics, Georgian Academy of Sciences,
Tbilisi, 380077, Georgia*

Abstract

We consider the problem of supersymmetry breaking in 5 dimensional $N = 1$ supersymmetric models with S^1/Z_2 compactification.

The remarkable success in the understanding of non-perturbative aspects of string theories gives a new insights into the particle phenomenology. One of the phenomenologically most promising approach has been proposed by Hořava and Witten within the 11-dimensional ($d=11$) supergravity compactified on S^1/Z_2 orbifold [1]. Below we study the problem of breaking of a rigid supersymmetry in $d = 5$ field-theoretic limit of the Hořava-Witten compactification. Consider the $d = 5$ off-shell $N = 1$ hypermultiplet $\mathcal{H} = (h^i, \psi, F^i)$ which consists of scalar field h^i and auxiliary field F^i , both being $SU(2)$ doublets and Dirac fermion $\psi = (\psi_L, \psi_R)^T$. These fields form two $d = 4$ $N = 1$ chiral multiplets $H_1 = (h^1, \psi_L, F^1)$ and $H_2 = (h^2, \psi_R, F^2)$ and are described by the $d = 5$ $N = 1$ SUSY Lagrangian:

$$\mathcal{L}_{hyper}^{(5)} = \left(\partial_M h^i\right)^+ \left(\partial^M h^i\right) + i\bar{\psi}\Gamma^M \partial_M \psi + \left(F^i\right)^+ \left(F^i\right). \quad (1)$$

Let us compactify the fifth coordinate x^4 on the orbifold S^1/Z_2 and define the transformation properties of fields entering in the hypermultiplet under the discrete Z_2 orbifold symmetry, which acts on the fifth coordinate as $x^4 \rightarrow -x^4$. A generic bosonic field transforms like $\varphi(x^m, x^4) = \mathcal{P}\varphi(x^m, -x^4)$ while the fermionic field transforms as $\eta(x^m, x^4) = \mathcal{P}i\sigma^3\eta(x^m, -x^4)$ where $\mathcal{P} = \pm 1$ is an intrinsic parity. Asserting $\mathcal{P} = 1$ to the components of multiplet H_1 and $\mathcal{P} = -1$ to the components of multiplet H_2 , one can see that the bulk Lagrangian is invariant under the action of the parity operator [2]. On the other hand at the orbifold fixed point $x^4 = 0$ one is left with $N=1$ supersymmetry acting on the $d = 4$ chiral multiplet H_1 with the modified auxiliary field $\tilde{F} = F^1 + \partial_4 h^2$.

*Talk at the 9th International Conference on Supersymmetry and Unification of Fundamental Interactions (SUSY01), Dubna, Russia, June 11-17, 2001

[†]E-mail: Masud.Chaichian@helsinki.fi

[‡]E-mail: Archil.Kobakhidze@helsinki.fi

[§]E-mail: tsulaia@thsun1.jinr.ru

Analogously, considering the $d = 5$ $N = 1$ vector multiplet which can be decomposed into $d = 4$ $N = 1$ vector $V = (A^m, \lambda_L^1, X^3)$ and chiral $\tilde{\Phi} = (\Sigma + iA^5, \lambda_L^2, X^1 + iX^2)$ multiplets and asserting $\mathcal{P} = 1$ for components of V and $\mathcal{P} = -1$ for components of $\tilde{\Phi}$, the bulk Lagrangian

$$\mathcal{L}_{Y.M.}^{(5)} = -\frac{1}{2g_5^2}F_{MN}^2 + \frac{1}{g_5^2} \left((D_M \Sigma)^2 + \bar{\lambda} i \Gamma^M D_M \lambda + (X^a)^2 - \bar{\lambda} [\Sigma, \lambda] \right) \quad (2)$$

is invariant under the action of the parity operator, while on the boundary $x^4 = 0$ we obtain $N = 1$ supersymmetry [3] realized on the vector multiplet V with $D = X^3 - \partial_4 \Sigma$.

The mechanism of supersymmetry breaking in these models [2] is based on a fact that any field configuration that breaks translational invariance and is not a BPS state breaks supersymmetry totally as well [4]. Such a stable non-BPS configurations with a purely finite gradient energy could appear in a compact spaces as well (or, more generally, in spaces with finite volume) if there exist moduli forming a continuous manifold of supersymmetric states. Consider first the case of the pure hypermultiplet in the $d = 5$ bulk. Besides the trivial vacuum configuration $\langle h^2 \rangle = 0$ there can exist non-trivial configuration

$$\langle h^2 \rangle = \epsilon x^4, \quad (3)$$

where ϵ is an arbitrary real constant. The configuration (3) is odd under the Z_2 orbifold transformation and breaks translational invariance in x^4 direction. However, the configuration (3) does not satisfy the ordinary periodicity condition on a S^1 circle, but rather the modified one,

$$h^2(x^m, x^4 + 2\pi R) = h^2(x^m, x^4) + 2\epsilon\pi R, \quad (4)$$

The Lagrangian density $\mathcal{L}_{hyper}^{(5)}$ remains single-valued and periodic. Thus, if we assume that h^2 and its superpartner ψ_R are defined modulo $2\epsilon\pi R$ on S^1/Z_2 then the configuration (3) will be perfectly compatible with S^1/Z_2 orbifold symmetries. The configuration (3) is stable (the variation of corresponding energy functional equals to zero) and spontaneously breaks $N = 1$ supersymmetry since $\langle F^1 + \partial_4 h^2 \rangle = \epsilon \neq 0$. Analogously there could be also x^4 -independent stable configuration

$$\langle h^1 \rangle = \epsilon r \sin \theta e^{i\varphi} \quad (5)$$

which breaks $N = 1$ supersymmetry on the boundary wall and is compatible with S^1/Z_2 orbifold symmetries.

The solution similar to (3) can be obtained as well for the Z_2 -odd scalar $\Sigma + iA_5$ in the case when vector supermultiplet lives in the bulk and non-trivial boundary condition analogous to (4) is assumed. Note also that the case of interacting gauge fields in the bulk actually reduces to the free theory considered above. Indeed the most general Lagrangian for supersymmetric gauge theories in $d = 5$ is expressed via the holomorphic prepotential being at most cubic [5] i.e., $\mathcal{F}(\Phi) = \frac{4\pi}{g_5^2}\Phi^2 + \frac{c}{3}\Phi^3$. However the S^1/Z_2 orbifold symmetry requires $c = 0$ and one is left again with the free theory described by (2).

Let us now consider the $N = 1$ chiral superfield $\Phi = (\phi, \chi_L, F_\Phi)$ localized on the $d = 4$ boundary $x^4 = 0$. The total Lagrangian has the form

$$\mathcal{L}_{hyper}^{(5)} + \left[\mathcal{L}_\Phi^{(4)} + \mathcal{L}_{\Phi H_1}^{(4)} \right] \delta(x^4), \quad (6)$$

where $\mathcal{L}_{\Phi H_1}^{(4)}(\Phi, H_1)$ describes the interactions between the chiral superfields $\Phi = (\phi, \chi_L, F_\Phi)$ and $H_1 = (h^1, \psi_L, F^1 + \partial_4 h^2)$ on the boundary $x^4 = 0$ through the superpotential $W_{\Phi H_1}(\Phi, H_1)$, while $\mathcal{L}_\Phi^{(4)}$ is the usual $d = 4$ Lagrangian for the chiral superfield with the superpotential W_Φ . Analyzing the equations of motion [2] one can see that if $\langle F^1 \rangle \neq 0$, and $\langle F_\Phi \rangle = 0$ then the degeneracy in $\langle h^2 \rangle$ is removed and for

$$\langle \frac{\partial W_{\Phi H_1}}{\partial h^1} \rangle = \alpha = \text{const} \quad (7)$$

we get the supersymmetry preserving configuration

$$\langle h^2 \rangle = -\alpha \theta(x^4). \quad (8)$$

Analogously adding to (2) the FI term $-2\eta (X^3 - \partial_4 \Sigma) \delta(x^4)$ one can find supersymmetry preserving configuration with

$$\langle X^3 \rangle = g_5^2 \eta \delta(x^4) \quad \text{and} \quad \langle \Sigma \rangle = g_5^2 \eta \theta(x^4). \quad (9)$$

Acknowledgments. The work of M.C. and A.K. was supported by the Academy of Finland under the Project 163394 and that of M.T. was supported by RFBR grants #99-02-18417 and #01-02-06531 and INTAS grant #00-00254.

References

- [1] P. Hořava and E. Witten, *Nucl. Phys.* **B460** (1996) 506; *Nucl. Phys.* **B475** (1996) 94.
- [2] M. Chaichian, A. Kobakhidze and M. Tsulaia, *Phys. Lett.* **B505** (2001) 222.
- [3] E. A. Mirabelli and M. Peskin, *Phys. Rev.* **D58** (1998) 065002.
- [4] G. Dvali and M. Shifman, *Nucl. Phys.* **B504** (1997) 127.
- [5] N. Seiberg, *Phys. Lett.* **B388** (1996) 753.